

Lung function tests in Asthma

Written by Dr Mike Levin (2008)
(Revised 2012)



(021) 830 5937

naepr@netactive.co.za

www.asthmasa.org

Asthma is a very common condition. It cannot be cured, but can be controlled with effective medication, good technique, and good monitoring by the patient and the doctor. Asthma is under control when patients are having no or few symptoms, no attacks needing hospital visits and no or little emergency reliever medication use. Another way of assessing asthma control is to do tests that measure the function of your lungs. These tests are sometimes also done to help with the diagnosis of asthma.

What are lung function tests?

Lung function tests are any test done that measures the functioning of the lungs. Some of these tests need sophisticated machines and highly trained staff, but some can be done by the patient at home.

Common lung function tests include:

- peak flow rates
- spirometry
- response and challenge tests

What is a peak flow test?

A peak flow test is a lung function test using a peak flow meter.

A peak flow meter is a small, simple machine that is held in the hand, and can be done at a doctor's office, clinic or even at home! The peak flow meter allows you to measure how fast the air can be blown out of your lungs. This is called the peak expiratory flow rate and is measured in litres per minute. A high peak flow rate means the airways are open, a low peak flow rate means the airways are closed, and the air cannot get out quickly. Peak flow tests are difficult or impossible to do in children below the age of four.

Why is it done?

Testing your peak expiratory flow may help:

- Diagnose asthma
- Guide the treatment and monitor the effectiveness of treatment for asthma.

- Measure how well your lungs are working to decide on the need for emergency reliever treatment such as reliever pumps, nebulisers and oral steroids.

How to use the peak flow meter:

1. You must be standing when using a peak flow meter.
2. Remove any food or chewing gum you may have in your mouth.
3. Set the pointer on the gauge of the peak flow meter to 0 (zero), and make sure your fingers are not in the way of the marker.
4. Attach the mouthpiece to the peak flow meter.
5. Hold the meter parallel with the floor.
6. Take a deep breath in, filling the lungs as much as possible.
7. Place the peak flow meter mouthpiece in your mouth and close your lips tightly around the outside of the mouthpiece. Do not put your tongue inside the mouthpiece.
8. Blow out as hard and as fast as you can. You do not need to breathe out for more than a second.
9. Read the value shown by the marker.
10. Reset the meter and blow into the peak flow meter 2 more times. Record your values each time you blow. If you cough or make a mistake during the testing, do the test again.
11. After you have blown into your peak flow meter 3 times, record the highest value on your daily record sheet.

Peak flow monitoring relies on your trying as hard as you can. For accurate results, it is important that you give the test your best effort every time.

What does it mean?

If there is a peak flow value of less than 80 % of expected, it means that the airways are narrowed. This can be measured with the peak flow meter before wheezing occurs, so it gives you the advantage of starting early treatment and preventing severe attacks.

Your peak flow readings will fall into different zones, the safe "green" zone (>80%), the orange "caution" zone (50% - 80%) and the red "danger" zone (<50%). Speak to your doctor about making a personal action plan that shows the peak flow values of each of these ranges for you, as well as a written action plan of what to do if your values fall in one of these zones. (See "Keeping asthma under control").

Signs of poor control include frequent low levels and a big difference between the morning values and evening values (of more than 15 – 20%). If this happens, your doctor may wish to improve your technique and frequency of taking controller medications, or may wish to add to or change your medications.

What else affects peak flow apart from asthma?

- Chronic obstructive airways disease,
- Time of day, recent use of broncodilator,
- Patient technique,
- False teeth,
- Incorrect posture,
- Type of meter.

What is spirometry?

Spirometry is a lung test that is done in a doctor's office or emergency room. If possible, it should be done on initial diagnosis of asthma, after treatment is started and symptoms have stabilized, and every year afterward. This test measures the flow of air out of your lungs using a forced breathing manoeuvre. This means the person takes as deep a breath as possible and then blows out as long and as hard as possible until the lungs are completely empty. The test is better than a peak flow rate, but cannot be done at home, and needs to be interpreted by a doctor or nurse who has experience in

doing the test. Your doctor or nurse may use it to assess the response of the lungs to treatment over time or to diagnose the type of lung disease that is present. This lung function test can show abnormalities that children and their parents may be unaware of. Spirometry is difficult to do in children below the age of six.

What are response and challenge tests?


Response tests measure lung function (using peak flow or spirometry) before any medication is given, and then again after a nebuliser or reliever asthma pump is given. One of the features of asthma, as opposed to other causes of wheezing, is that asthma will respond to the medicine with an improvement in the lung function tests of more than 15%. This is sometimes called a reversibility test or a post bronchodilation test.

Challenge tests are done less frequently than response tests. A common challenge test is the exercise challenge. Children who are suspected of having asthma are asked to perform lung function tests as a baseline. Then they exercise for 6 minutes until their heart rates go up to over 170. The lung function tests are repeated after a recovery period of 5 minutes. Many asthmatics will show a drop in their lung function tests because the airways have closed after the exercise. If this happens a response test should immediately be done to prove a good response to the asthma medication.

In specialized centres chemicals or allergens may be used as a challenge instead of exercise to cause tightening of the airways. These are done very rarely and only under expert supervision.

What is exhaled Nitric Oxide?

This is a breathing test to see how much inflammation (and swelling) is in your lungs. It can only be done at a doctor's office and is only available at a few places in South Africa. To do the test, you breathe slowly and constantly into a mouthpiece which measures the nitric oxide gas given off by inflamed lungs! The test is especially useful if your doctor is not sure whether your lung problem is asthma or not, and can also be used to measure the response to therapy, and therefore control of your asthma. If the measurement is high, then, lots of inflammation is present, and medication, technique and the regularity of using controller medication needs to be reassessed.



Join the NAEP!
sign up on www.asthmasa.org